

Class 2

Densities and Distribution Functions

A continuous random variable X has a **density** $f(x) \geq 0$ that integrates to one, and probabilities are areas under it,

$$P(a < X \leq b) = \int_a^b f(x) dx = F(b) - F(a),$$

where $F(x) = P(X \leq x)$ is the **CDF**, with $F'(x) = f(x)$. The density is not a probability: it can exceed one, and any single value has probability zero. The right tail, or **survival function**, is $P(X > x) = 1 - F(x)$. Sums become integrals, so $E(g(X)) = \int g(x)f(x) dx$, and everything we did with discrete random variables carries over.

If $Y = g(X)$ with g strictly increasing, then $P(Y \leq y) = P(X \leq g^{-1}(y))$, and differentiating gives

$$f_Y(y) = f_X(g^{-1}(y)) \frac{dg^{-1}(y)}{dy}. \quad (1)$$

The derivative accounts for how the transformation stretches or compresses intervals.

Uniform and Exponential

The **uniform** distribution on $[a, b]$ has constant density $1/(b - a)$, mean $(a + b)/2$ and variance $(b - a)^2/12$. It drives simulation through the **inverse transform method**: if F is continuous and strictly increasing and $U \sim \text{Uniform}(0, 1)$, then $X = F^{-1}(U)$ has CDF F .

The **exponential** distribution $T \sim \text{Exp}(\lambda)$ models the waiting time for an event that occurs at a constant rate λ , such as a default with constant intensity,

$$f(t) = \lambda e^{-\lambda t}, \quad P(T > t) = e^{-\lambda t}, \quad E(T) = \frac{1}{\lambda}, \quad V(T) = \frac{1}{\lambda^2}.$$

Solving $u = 1 - e^{-\lambda t}$ gives the simulation recipe $T = -\ln(U)/\lambda$.

The Gamma Distribution

The exponential distribution describes the waiting time for one event. The gamma distribution describes, among other things, the waiting time for several. Its density involves the **gamma function**

$$\Gamma(\alpha) = \int_0^{\infty} x^{\alpha-1} e^{-x} dx, \quad \alpha > 0. \quad (2)$$

Integrating by parts gives $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$, and since $\Gamma(1) = 1$, $\Gamma(n) = (n - 1)!$ for every positive integer n : the gamma function extends the factorial to non-integer arguments. We will also need $\Gamma(1/2) = \sqrt{\pi}$.

A random variable $X \sim \text{Gamma}(\alpha, \lambda)$, with shape $\alpha > 0$ and rate $\lambda > 0$, has density

$$f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x}, \quad x > 0.$$

The substitution $u = \lambda x$ in (2) shows that it integrates to one. For $k \geq 1$, the integrand of $E(X^k)$ is proportional to a gamma density with shape $\alpha + k$, so

$$E(X^k) = \frac{\Gamma(\alpha + k)}{\Gamma(\alpha)\lambda^k}, \quad E(X) = \frac{\alpha}{\lambda}, \quad V(X) = \frac{\alpha}{\lambda^2}. \quad (3)$$

For $\alpha \leq 1$ the density decreases from its value at zero, while for $\alpha > 1$ it rises from zero to a peak at $(\alpha - 1)/\lambda$ and then decays. With $\alpha = 1$ the gamma is the exponential distribution. A sum of n independent $\text{Exp}(\lambda)$ waiting times is $\text{Gamma}(n, \lambda)$, so the time of the n -th event in a Poisson process is gamma distributed.

The Normal Distribution

$X \sim \mathcal{N}(\mu, \sigma^2)$ has density $e^{-(x-\mu)^2/(2\sigma^2)}/\sqrt{2\pi\sigma^2}$, mean μ and variance σ^2 . Every normal probability reduces to the standard normal CDF Φ through the Z-score,

$$P(X \leq x) = \Phi\left(\frac{x - \mu}{\sigma}\right), \quad \Phi(-z) = 1 - \Phi(z). \quad (4)$$

With z_α defined by $P(Z > z_\alpha) = \alpha$ ($z_{0.05} = 1.64$, $z_{0.025} = 1.96$, $z_{0.005} = 2.58$), the $(1 - \alpha)$ **probability interval** of X is $[\mu - z_{\alpha/2}\sigma, \mu + z_{\alpha/2}\sigma]$. It says where X is likely to fall; it is not a confidence interval.

If $Z \sim \mathcal{N}(0, 1)$, then $Z^2 \sim \text{Gamma}(1/2, 1/2)$, the **chi-squared** distribution with one degree of freedom, so $E(Z^2) = 1$, $V(Z^2) = 2$ and $E(Z^4) = 3$. A sum of k independent squared standard normals is chi-squared with k degrees of freedom, $\text{Gamma}(k/2, 1/2)$, with mean k and variance $2k$.

The Lognormal Distribution

If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Y = e^X$ is **lognormal**: positive, skewed to the right, and the standard model for a stock price. Here μ and σ^2 are the mean and variance of $\ln(Y)$, not of Y . Probabilities come from the logarithm, $P(Y \leq y) = \Phi((\ln(y) - \mu)/\sigma)$, and the moments are

$$E(Y^\alpha) = e^{\alpha\mu + \frac{1}{2}\alpha^2\sigma^2}, \quad E(Y) = e^{\mu + \frac{1}{2}\sigma^2}, \quad \text{SD}(Y) = E(Y)\sqrt{e^{\sigma^2} - 1}. \quad (5)$$

The median is e^μ , below the mean, because the long right tail pulls the mean up (Figure 1). The $(1 - \alpha)$ probability interval $[e^{\mu - z_{\alpha/2}\sigma}, e^{\mu + z_{\alpha/2}\sigma}]$ divides and multiplies the median by the same factor, so it is not symmetric.




Figure 1: Density of a stock price in one year, $Y = 100e^X$ with $X \sim \mathcal{N}(0.05, 0.30^2)$. The median $100e^{0.05}$ lies below the mean $100e^{0.05+0.045}$.

The Bivariate Normal Distribution

Correlated normal returns are built from independent standard normals Z_1 and Z_2 ,

$$X = \mu_X + \sigma_X Z_1, \quad Y = \mu_Y + \sigma_Y (\rho Z_1 + \sqrt{1 - \rho^2} Z_2), \quad (6)$$

which gives $\text{Cov}(X, Y) = \rho \sigma_X \sigma_Y$. Three properties follow from the construction.

1. **Linear combinations.** $aX + bY$ is normal with mean $a\mu_X + b\mu_Y$ and variance $a^2\sigma_X^2 + b^2\sigma_Y^2 + 2ab\rho\sigma_X\sigma_Y$.
2. **Conditional distribution.** Given $X = x$, Y is normal with mean $\mu_Y + \rho(\sigma_Y/\sigma_X)(x - \mu_X)$ and variance $\sigma_Y^2(1 - \rho^2)$. The conditional mean is the regression line, with slope $\text{Cov}(X, Y)/V(X)$.
3. **Independence.** X and Y are independent if and only if $\rho = 0$. This is special to jointly normal variables.

Heavy Tails

Daily returns have far more extreme moves than the normal distribution predicts. The **Student t** distribution with ν degrees of freedom, $T = Z/\sqrt{V/\nu}$ with V chi-squared and independent of Z , is normal given a random scale. Its density decays like the power $|t|^{-(\nu+1)}$, so $E(|T|^k)$ is finite only for $k < \nu$, the variance $\nu/(\nu - 2)$ requires $\nu > 2$, and $E(e^{\theta T}) = \infty$ for every $\theta \neq 0$. As $\nu \rightarrow \infty$ it approaches the standard normal.

Practice Problems

1. The recovery rate R on a defaulted bond, per dollar of face value, is uniformly distributed on $[0.2, 0.6]$.
 - a. Compute $E(R)$ and $SD(R)$, and the probability that $R > 0.5$.
 - b. A contract pays $\max(R - 0.3, 0)$ per dollar of face value. Compute its expected payoff.
 - c. A computer generates the uniform draws $u = 0.10$ and $u = 0.75$. Use the inverse transform method to turn them into simulated recovery rates.
2. The time W , in minutes, until the second trade in an illiquid bond has a Gamma($2, \lambda$) distribution with $\lambda = 2$ per minute.
 - a. Compute $E(W)$, $SD(W)$ and the value at which the density of W peaks.
 - b. Integrate the density by parts to show that $P(W \leq t) = 1 - e^{-\lambda t}(1 + \lambda t)$, and compute $P(W \leq 1)$.
 - c. Let $D = 60W$ be the same waiting time measured in seconds. Use (1) to find its density f_D , and show that D is again gamma distributed. Which parameter changes, and why?
3. The daily return of a \$10 million portfolio is normal with mean 0.0004 and standard deviation 0.015.
 - a. What is the probability that the portfolio loses more than 2% in a day?
 - b. Using $z_{0.05} = 1.64$, find the dollar loss that is exceeded on only 5% of days.
 - c. A risk manager estimates the daily variance with $\hat{\sigma}^2 = \frac{1}{21} \sum_{t=1}^{21} (R_t - \mu)^2$, using 21 independent daily returns R_t and the true mean μ . Use the chi-squared distribution to compute $E(\hat{\sigma}^2)$ and the ratio $SD(\hat{\sigma}^2)/\sigma^2$.
4. A stock trades at $S_0 = 50$. Its log return over the year is $\mathcal{N}(\mu, \sigma^2)$ with $\mu = 0.06$ and $\sigma = 0.30$.
 - a. What is the probability that a call option with strike 60 and one year to maturity finishes in the money, that is, $P(S_1 > 60)$?
 - b. Compute $E(S_1)$ and compare it with the median price $S_0 e^\mu$.